

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau






INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 235/06, 401/06, 401/14, 209/08, 403/06, A61K 31/445, 31/40, 31/41		A1	(11) International Publication Number: WO 99/61426
			(43) International Publication Date: 2 December 1999 (02.12.99)
(21) International Application Number: PCT/US99/11222		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 21 May 1999 (21.05.99)		Published With international search report.	
(30) Priority Data: 60/086,531 22 May 1998 (22.05.98) US 09/128,137 3 August 1998 (03.08.98) US 09/275,176 24 March 1999 (24.03.99) US			
(71) Applicant: SCIOS, INC. [US/US]; 2450 Bayshore Parkway, Mountain View, CA 94403 (US).			
(72) Inventors: MAVUNKEL, Babu, J.; 1348 Avoset Terrace, Sunnyvale, CA 94087 (US). LIU, David, Y.; 201 Ferne Avenue, Palo Alto, CA 94306 (US). SCHREINER, George, F.; 12774 Leander Drive, Los Altos Hills, CA 94022 (US). LEWICKI, John, A.; 308 Escobar Avenue, Los Gatos, CA 95030 (US). PERUMATTAM, John, J.; 30 Chester Circle, Los Altos, CA 94022 (US).			
(74) Agents: MURASHIGE, Kate, H. et al.; Morrison & Foerster LLP, 2000 Pennsylvania Avenue, N.W., Washington, DC 20006-1888 (US).			

(54) Title: HETEROCYCLIC COMPOUNDS AND METHODS TO TREAT CARDIAC FAILURE AND OTHER DISORDERS

(57) Abstract

Compounds of formulae α or β , and the pharmaceutically acceptable salts thereof, wherein each of Z^1 and Z^2 is independently CR^4 or N; where each R^4 is independently H or is alkyl (1-6C) or aryl, each of said alkyl or aryl optionally including one or more heteroatoms selected from O, S and N and optionally substituted by one or more of halo, OR, SR, NR_2 , RCO, COOR, $CONR_2$, OOCR, or NROCR where R is H or alkyl (1-6C), or by one or more CN or =O, or by one or more aliphatic or aromatic 5- or 6-membered rings optionally containing 1-2 heteroatoms; R^1 is formula (I); wherein X^1 is CO or an isostere thereof; m is 0 or 1; Y is optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl or two Y taken together may form an alkylene (2-3C) bridge; n is 0 or 2; Z^3 is CH or N; X^2 is CH, CH_2 or an isostere thereof; and Ar consists of one or two phenyl moieties directly coupled to X^2 optionally substituted by halo, nitro, alkyl(1-6C), alkenyl(1-6C), alkynyl(1-6C), CN or CF_3 , or by RCO, COOR, $CONR_2$, NR_2 , OR, SR, OOCR or NROCR wherein R is H or alkyl(1-6C) or by phenyl, itself optionally substituted by the foregoing substituents; R^2 is H, or is alkyl(1-6C) or aryl, each or said alkyl or aryl optionally including one heteroatom which is O, S or N, and optionally substituted by one or more of halo, OR, SR, NR_2 , RCO, COOR, $CONR_2$, OOCR, or NROCR where R is H or alkyl(1-6C), alkynyl(1-6C), or by one or more CN or =O, or by one or more aliphatic or aromatic 5- or 6-membered rings optionally containing 1-2 heteroatoms; R^3 is H, halo, NO_2 , alkyl(1-6C), alkenyl(1-6C), alkynyl(1-6C), CN, OR, SR, NR_2 , RCO, COOR, $CONR_2$, OOCR, or NROCR where R is H or alkyl(1-6C) are disclosed. These compounds are selective inhibitors of p38 α kinase.



Atty. Docket No. 3015/6/US
Serial No. 10/021,780
Anantanarayan et al.
Reference 44 of 77

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

HETEROCYCLIC COMPOUNDS AND METHODS TO TREAT CARDIAC FAILURE AND OTHER DISORDERS

5 Technical Field

The invention is directed to compounds that are useful in treating inflammation and that contain piperazine or piperidine moieties coupled to the 5- or 6-position of indole, benzimidazole or benzotriazole. More particularly, the invention concerns novel ortho substituted indoles and N-substituted indoles as well as methods
10 to treat heart and kidney conditions using these compounds and derivatives thereof.

Background Art

A large number of chronic and acute conditions have been recognized to be associated with perturbation of the inflammatory response. A large number of
15 cytokines participate in this response, including IL-1, IL-6, IL-8 and TNF. It appears that the activity of these cytokines in the regulation of inflammation rely at least in part on the activation of an enzyme on the cell signaling pathway, a member of the MAP kinase family generally known as p38 and alternatively known as CSBP and RK. This kinase is activated by dual phosphorylation after stimulation by
20 physiochemical stress, treatment with lipopolysaccharides or with proinflammatory cytokines such as IL-1 and TNF. Therefore, inhibitors of the kinase activity of p38 are useful antiinflammatory agents.

PCT applications WO98/28292, WO98/06715, WO98/07425, and WO
96/40143, all of which are incorporated herein by reference, describe the relationship
25 of p38 kinase inhibitors with various disease states. As mentioned in these applications, inhibitors of p38 kinase are useful in treating a variety of diseases associated with chronic inflammation. These applications list rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis and other arthritic conditions, sepsis, septic shock, endotoxic shock, Gram-negative sepsis, toxic shock syndrome,
30 asthma, adult respiratory distress syndrome, stroke, reperfusion injury, CNS injuries such as neural trauma and ischemia, psoriasis, restenosis, cerebral malaria, chronic


pulmonary inflammatory disease, silicosis, pulmonary sarcosis, bone resorption diseases such as osteoporosis, graft-versus-host reaction, Crohn's Disease, ulcerative colitis including inflammatory bowel disease (IBD) and pyresis.

The above-referenced PCT applications disclose compounds which are p38
 5 kinase inhibitors said to be useful in treating these disease states. These compounds are either imidazoles or are indoles substituted at the 3- or 4-position with a piperazine or piperidine ring linked through a carboxamide linkage. Additional compounds which are conjugates of piperazines with indoles are described as insecticides in WO97/26252, also incorporated herein by reference.


10

Disclosure of the Invention

The invention is directed to compounds useful in treating inflammation generally, including specific conditions such as those described in the Background section above. Certain novel compounds have been found to inhibit p38 kinase, in
 15 particular, p38 kinase α and are thus useful in treating diseases mediated by this enzyme. The compounds of the invention are of the formula:



preferably those of the formulas:




and the pharmaceutically acceptable salts thereof,

wherein each of Z^1 and Z^2 is independently CR^4 or N;

where each R^4 is independently H or is alkyl (1-6C) or aryl, each of said alkyl or aryl optionally including one or more heteroatoms selected from O, S and N and optionally substituted by one or more of halo, OR, SR, NR_2 , RCO, COOR, $CONR_2$,
 5 OOCR, or NROCR where R is H or alkyl (1-6C), or by one or more CN or =O, or by one or more aliphatic or aromatic 5- or 6-membered rings optionally containing 1-2 heteroatoms;

R^1 is



10 wherein

X^1 is CO or an isostere thereof;

m is 0 or 1;

Y is optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl or two Y taken together may form an alkylene (2-3C) bridge;

15 n is 0, 1 or 2;

Z^3 is CH or N;

X^2 is CH, CH_2 or an isostere thereof; and

Ar consists of one or two phenyl moieties directly coupled to X^2 optionally substituted by halo, nitro, alkyl (1-6C), alkenyl (1-6C), alkynyl (1-6C), CN or CF_3 , or
 20 by RCO, COOR, $CONR_2$, NR_2 , OR, SR, OOCR or NROCR wherein R is H or alkyl (1-6C)

or by phenyl, itself optionally substituted by the foregoing substituents;

R^2 is H, or is alkyl (1-6C) or aryl each of said alkyl or aryl optionally including one or more heteroatoms which are O, S or N, and optionally substituted by one or
 25 more of halo, OR, SR, NR_2 , RCO, COOR, $CONR_2$, OOCR, or NROCR where R is H or alkyl (1-6C), or by one or more CN or =O, or by one or more aliphatic or aromatic 5- or 6-membered rings optionally containing 1-2 heteroatoms;

R^3 is H, halo, NO_2 , alkyl (1-6C), alkenyl (1-6C), alkynyl (1-6C), CN, OR, SR, NR_2 , RCO, COOR, $CONR_2$, OOCR, or NROCR where R is H or alkyl (1-6C).

Thus, in one aspect, the invention is directed to compounds of the formulas set forth above. In other aspects, the invention is directed to methods to produce these compounds, to pharmaceutical compositions containing them, and to methods of treating inflammation using these compounds. The invention is also directed to
5 treating conditions associated with cardiac failure using the invention compounds and other compounds described herein.

Modes of Carrying Out the Invention

The compounds of formulas 1-4 are useful in a variety of physiological
10 contexts, as further described below. Preferred embodiments include those wherein both Z^1 and Z^2 are CH or where Z^1 is CR^4 and Z^2 is CH; thus, among the preferred compounds of the invention are derivatives of indole. Especially preferred substituents at the 3-position are those coupled through carboxamide linkages. Thus, some preferred embodiments of R^4 are of the formula $RNHCO-$ wherein R is alkyl or
15 substituted alkyl.

In general, substituents on the nitrogen-containing portion of the indole, benzimidazole, or benztriazole nucleus are designed to enhance solubility. Thus, typically, the substituents R^2 and R^4 are polar or contain polar groups.

In other preferred embodiments, the substituents shown for the compounds of
20 the invention are as set forth below.

In regard to R^1 :

X^1 is CO or an isostere thereof. Thus, in addition to CO, X^1 may be CH_2 , SO, SO_2 , or CHOH. CO is preferred.

Z^3 is CH or N; $Z^3 = CH$ is preferred.

25 Typically m is 1; however, in some compounds of the invention, m can be 0; thus, this substituent is a five-membered ring.

X^2 is CH_2 if Ar consists of a single phenyl moiety or CH if Ar consists of two phenyl moieties or may be an isostere thereof. Thus, for appropriate embodiments of Ar, X^2 may be any of the alternatives set forth above for X^1 .


30 The phenyl moieties represented by Ar may optionally be substituted by substituents including alkyl (1-6C), halo, RCO, COOR, $CONR_2$, OR, SR, NR_2 ,

5 OOCR, NROCR, NO₂, CN, or CF₃, wherein R is H or alkyl (1-6C). The phenyl moieties may also be substituted with an additional phenyl residue, preferably at the 4-position. The additional phenyl residue may itself be substituted with the substituents set forth above. The additional phenyl may be substituted in all five positions, but preferably less, preferably in 1-2 positions or not at all. Preferred substituents include alkyl (1-6C), OR, NR₂ and halo, especially halo and OCH₃. The substituents may occupy all five positions of the phenyl substituent, preferably 1-2 positions or the phenyl may be unsubstituted.

10 n may be 0, 1 or 2, and is preferably 0. However, when n is 1, Y is present and may be alkyl, arylalkyl or aryl, all of which may optionally be substituted by the substituents set forth above with regard to Ar. When n is 2, both Y groups together may constitute an alkylene bridge. A preferred bridge is an ethylene bridge. Preferred embodiments of Y when n is 1 include unsubstituted alkyl and unsubstituted arylalkyl.

15 With regard to R²:

R² is preferably H, but may also be a suitable substituent. Such substituents are typically and preferably alkyl or substituted alkyl. The alkyl or substituted alkyl may optionally include one or more heteroatoms which can be O, N or S, preferably N and O. Permitted substitutions on the alkyl group are set forth above; preferred substituents include OR, where R is H or alkyl (1-6C) and =O. Also included among the preferred substituents on the alkyl group are cyclic moieties, such as piperazine, pyridine, piperidine, phenyl, and the like. Preferably, the alkyl embodiments of R² contain 0, 1 or 2 substituents. Among preferred embodiments of R² are included those of the formula -(CO)O-Y' wherein Y' is, for example, -(CH₂)_nNR₂, where n is an integer of 0-6 and R is as defined above; or Y' is, for example, an aliphatic or aromatic ring system, such as



Additional illustrative embodiments of R^2 include nicotinoyl and its isomers, acryloyl, and substituents of the general formula $Y'(CH_2)_nNH(CH_2)_nCHOH(CH_2)_n-$ wherein Y' is a generic substituent such as optionally substituted alkyl, piperazinyl, piperidinyl, cyclohexyl, phenyl or methoxy, and the like and wherein each n is

5 independently an integer of 1-3. Y' is quite variable and can generally include any noninterfering moiety. Additional embodiments include those of the general formula $Y'NH(CH_2)_n-CO$, wherein Y' and n are as described above; also included are those of the general formula $Y'(CH_2)_nNH(CH_2)_nCO$ where Y' and n are as described above; and those of the formula $Y'(CH_2)_nCO$ and $Y'(CH_2)_nNHCO$, wherein Y' and n are as defined


10 above; and those of the formula $R_2N(CH_2)_n-$ wherein R is alkyl (1-6C) and n is an integer of 1-3.

With respect to R^3 :

Although R^3 may be H, other embodiments are included and may be preferred. These include halo, OR, NR_2 , and alkyl (1-6C), as particularly desirable.

15 In embodiments wherein Z^1 or Z^2 , preferably Z^1 , is CR^4 , where R^4 is other than H, preferred embodiments of R^4 include those of the formula $R_2N(CH_2)_n-$ wherein each R is independently alkyl (1-6C) or H and n is an integer of 1-6; or of the formula $Y'(-CH_2)_n-$ wherein Y' is as defined above and n is an integer of 1-6; or those of the formula $Y'NHCO$; or those of the formula R_2NCO , wherein the R_2 substituents taken

20 together form a ring which may itself be substituted, preferably by alkyl, arylalkyl, and the like. When R^4 is $Y'(CH_2)_n-$, for example, Y' may be




Additional illustrative embodiments of R^4 include 2-, 3- and 4-pyridyl, 2-, 3- and 4-piperidyl.

25

The compounds of formulas (1)-(4) may be supplied in the form of their pharmaceutically acceptable acid-addition salts including salts of inorganic acids such

as hydrochloric, sulfuric, hydrobromic, or phosphoric acid or salts of organic acids such as acetic, tartaric, succinic, benzoic, salicylic, and the like. If a carboxyl moiety is present, these compounds may also be supplied as a salt with a pharmaceutically acceptable base, including inorganic bases such as sodium hydroxide, potassium hydroxide, calcium hydroxide, ammonium hydroxide and the like or a salt with an organic base such as caffeine.

Particularly preferred compounds of the invention are of formulas (5) and (6):



In these compounds, R^1 is of the formula shown, wherein each X^3 is independently halo, alkyl (1-6C), OR or NR_2 , wherein R is H or alkyl (1-6C), and p is an integer of 0-3. R^2 , R^3 and R^4 are as defined above.

Also preferred are similar compounds where the positions of R^3 and the illustrated embodiment of R^1 are reversed; i.e., R^3 is at position 5 and R^1 is in position 6.

15 Synthesis of the Invention Compounds

The compounds of the invention can be synthesized by a variety of methods most of them known in the art per se. The indole, benzimidazole, or benotriazole moiety may be supplied per se and the substituent R^1 coupled thereto. R^1 may be supplied as such, or its synthesis may be completed when the piperazyl or piperidyl residue is already coupled to the indole, benzimidazole or benotriazole moiety.

20 Alternatively, especially in embodiments wherein R^3 represents a non-hydrogen substituent, the appropriately substituted p-aminobenzoic acid derivative may be cyclized and then substituted with piperazine or piperidine.


Thus, for example, as shown in Reaction Scheme 1, a piperazine protected with tert-butyloxycarbonyl (BOC) is coupled to 5-carboxybenzimidazole (or 5-

25

carboxy-indole, or 5-carboxy-benzotriazole) in a reaction mixture containing a coupling agent such as EDAC in an inert, aprotic solvent to obtain the coupled carboxamide which is then deprotected and treated with substituted or unsubstituted benzyl halides or benzoyl halides.

5

Scheme 1




R^a = for example, 2,6-difluorophenyl; 3,4-difluorophenyl; 2,3-difluorophenyl; 3,5-difluorophenyl; 3-chlorophenyl; 4-chlorophenyl; 4-carboxymethylphenyl; 4-methoxyphenyl; 4-trifluoromethoxyphenyl; 4-methylphenyl; 6-chloropiperonyl; t-butylcarboxyphenyl; 3-trifluorophenyl; 2,4-dichlorophenyl; 3,4-dichlorophenyl; phenyl; methoxyphenyl; or p-toluy.

10

15


Alternatively, as shown in Reaction Scheme 2, 5-carboxylated benzimidazole (or indole or benzotriazole) is reacted with a piperazine or piperidine moiety already substituted by X²-Ar. In this reaction, the piperazyl or piperidyl derivative is directly reacted with the carboxylated bicycloheteroatom-containing nucleus in the presence of a coupling agent such as EDAC in the presence of an inert solvent as set forth above.

Scheme 2



In order to form the substituted piperazine required for Scheme 2, piperazine is first converted to the BOC derivative and then reacted with ArCHO in the presence of a borohydride under acidic conditions to give the substituted piperazine as shown in Reaction Scheme 3.


Scheme 3



An alternative for coupling derivatized piperazine or piperidine to indole, benzimidazole or benzotriazole is shown in Reaction Scheme 4. In this reaction, the

piperazine or piperidine ring is derivatized to a suitable leaving group as shown and then treated with a base such as NaH in an inert solvent to obtain the desired conjugate.

Scheme 4




5

Another alternative is shown in Reaction Scheme 5. In this approach, a protected piperidone is reacted in the presence of a base, such as NaH, with the appropriate phosphonate ester to obtain a protected benzylene piperidine. The product is then deprotected and reacted with the carboxylate of indole, benzimidazole or benzotriazole using an appropriate dehydrating agent. The product is then reduced to the desired arylalkylated piperidine derivative.

1.0


Scheme 5



- Reaction Scheme 6 illustrates a method for preparing compounds of the invention in which the indole is substituted in the 6-membered ring thereof. In
- 5 Reaction Scheme 6, the appropriately substituted aniline is reacted with 1-methylmercaptyl-2,2-dialkoxyethane in the presence of tertiary butyryl chloride and base to provide the desired indole. Depending on the nature of the substitution of the aniline starting material, more than one isomer may result as shown. The methylmercaptyl group remaining on the 5-membered ring is reduced with Raney
- 10 nickel and a mandatory methyl group included on the original aniline moiety is hydrolyzed to the corresponding carboxylic acid. The resulting acid is then reacted with the desired piperidine or piperazine derivative in the presence of a coupling agent such as EDC.


- 12 -

Scheme 6

X = Cl, OCH_3 , CH_3


- 5 Alkylation of the nitrogens on the indole, benzimidazole or benzotriazole nucleus in the compounds *per se* is carried out by conventional means by reacting the halide of the substituent to be added in the presence of base and acetone, as shown in the illustrative alternative depictions of Scheme 7.

Scheme 7



10


or



where X = H, OMe, Cl;

each R is H or alkyl;

n is an integer; or




5

where X = H, OHC₃, Cl, CH₃, etc.;

each R is H, alkyl, aryl


or together both R form



Substituents at the 3-position of indole can be modified using the general procedures shown in Scheme 8:

5

Scheme 8



where X = OMe, Cl, CH₃;

each R is H, alkyl, aryl or together the R groups are piperazinyl,

4-benzylpiperazinyl, etc.

10 For synthesis of compounds wherein n is 1 -- i.e., wherein the piperidine ring contains one additional substituent other than those mandated in the compounds of the invention, the 4-substituted piperidine is first protected using BOC_2O in THF or other aprotic solvent and then reacted with, for example, an alkyl iodide in the presence of S-butyl lithium/TMEDA using, for example, ether as a solvent to produce the

15 alkylated piperidine. The alkylated piperidine is then converted to the invention compound by deprotection followed by formation of the carboxamido linkage to the indoyl residue. This is exemplified below.

For compounds of the invention that are indoles substituted at the 3-position, the Reaction Scheme shown at the beginning of Example 23 may conveniently be

20 used. Typically, the carboxamide starting material is treated with trifluoroacetic anhydride to obtain the trifluoroacetyl intermediate, which is also a compound of the invention. Upon treatment with base, the 3-carboxylic acid is formed which can then be reacted with a suitable amine to obtain additional compounds of the invention.

Administration and Use

The compounds of the invention are useful in treating conditions associated with inflammation. Thus, the compounds of formulas (1)-(4) or their pharmaceutically acceptable salts are used in the manufacture of a medicament for prophylactic or therapeutic treatment of mammals, including humans, in respect of conditions characterized by excessive production of cytokines and/or inappropriate or unregulated cytokine activity on such cells as cardiomyocytes, cardiofibroblasts and macrophages.

The compounds of the invention inhibit the production of cytokines such as TNF, IL-1, IL-6 and IL-8, cytokines that are important proinflammatory constituents in many different disease states and syndromes. Thus, inhibition of these cytokines has benefit in controlling and mitigating many diseases. The compounds of the invention are shown herein to inhibit a member of the MAP kinase family variously called p38 MAPK (or p38), CSBP, or SAPK-2. The activation of this protein has been shown to regulate the production of prostanoids, such as PGE₂, and matrix metalloproteinases, such as collagenase-3, and to accompany exacerbation of the diseases in response to stress caused, for example, by treatment with lipopolysaccharides or cytokines such as TNF and IL-1. Inhibition of p38 activity, therefore, is predictive of the ability of a medicament to provide a beneficial effect in treating diseases such as coronary artery disease, congestive heart failure, cardiomyopathy, myocarditis, vasculitis, restenosis, such as occurs following coronary angioplasty, atherosclerosis, rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis and other arthritic conditions, multiple sclerosis, acute respiratory distress syndrome (ARDS), asthma, chronic obstructive pulmonary disease (COPD), silicosis, pulmonary sarcosis, sepsis, septic shock, endotoxic shock, toxic shock syndrome, heart and brain failure (stroke) that are characterized by ischemia and reperfusion injury, surgical procedures, such as transplantation procedures and graft rejections, cardiopulmonary bypass, coronary artery bypass graft, CNS injuries, including open and closed head trauma, inflammatory eye conditions such as conjunctivitis and uveitis, acute renal failure, glomerulonephritis, inflammatory bowel diseases, such as Crohn's disease or ulcerative colitis, graft vs. host disease, bone

resorption diseases like osteoporosis, type II diabetes, pyresis, psoriasis, cachexia, viral diseases such as those caused by HIV, CMV, Herpes, and cerebral malaria, tumor metastases and acute pain, such as that accompanying dental surgery, dysmenorrhea and post-orthopedic surgery.

5 Within the last several years, p38 has been shown to comprise a group of MAP kinases designated p38 α , p38 β , p38 γ and p38 δ . Jiang, Y. *et al. J Biol Chem* (1996) 271:17920-17926 first reported characterization of p38 β as a 372-amino acid protein closely related to p38 α . Kumar, S. *et al. Biochem Biophys Res Comm* (1997) 235:533-538 and Stein, B. *et al. J Biol Chem* (1997) 272:19509-19517 reported a
10 second isoform of p38 β , p38 β 2, containing 364 amino acids with 73% identity to p38 α . All of these reports show evidence that p38 β is activated by proinflammatory cytokines and environmental stress, although the second reported p38 β isoform, p38 β 2, appears to be preferentially expressed in the CNS, heart and skeletal muscle compared to the more ubiquitous tissue expression of p38 α . Furthermore, activated
15 transcription factor-2 (ATF-2) was observed to be a better substrate for p38 β 2 than for p38 α , thus suggesting that separate mechanisms of action may be associated with these forms. The physiological role of p38 β 1 has been called into question by the latter two reports since it cannot be found in human tissue and does not exhibit appreciable kinase activity with the substrates of p38 α .


20 The identification of p38 γ was reported by Li, Z. *et al. Biochem Biophys Res Comm* (1996) 228:334-340 and of p38 δ by Wang, X., *et al., J Biol Chem* (1997) 272:23668-23674 and by Kumar, S., *et al., Biochem Biophys Res Comm* (1997) 235:533-538. The data suggest that these two p38 isoforms (γ and δ) represent a unique subset of the MAPK family based on their tissue expression patterns, substrate
25 utilization, response to direct and indirect stimuli, and susceptibility to kinase inhibitors.

 Various results with regard to response to drugs targeting the p38 family as between p38 α and either the putative p38 β 1 or p38 β 2 or both were reported by Jiang, Kumar, and Stein cited above as well as by Eyers, P.A. *et al. Chem and Biol* (1995)
30 5:321-328. An additional paper by Wang, Y. *et al. J Biol Chem* (1998) 273:2161-2168 suggests the significance of such differential effects. As pointed out by Wang, a


number of stimuli, such as myocardial infarction, hypertension, valvular diseases, viral myocarditis, and dilated cardiomyopathy lead to an increase in cardiac workload and elevated mechanical stress on cardiomyocytes. These are said to lead to an adaptive hypertrophic response which, if not controlled, has decidedly negative consequences. Wang cites previous studies which have shown that in ischemia reperfusion treated hearts, p38 MAPK activities are elevated in association with hypertrophy and programmed cell death. Wang shows in the cited paper that activation of p38 β activity results in hypertrophy, whereas activation of p38 α activity leads to myocyte apoptosis. Thus, selective inhibition of p38 α activity as compared to p38 β activity will be of benefit in treating conditions associated with cardiac failure. These conditions include congestive heart failure, cardiomyopathy, myocarditis, vasculitis, vascular restenosis, valvular disease, conditions associated with cardiopulmonary bypass, coronary artery bypass, grafts and vascular grafts. Further, to the extent that the α -isoform is toxic in other muscle cell types, α -selective inhibitors would be useful for conditions associated with cachexia attributed to TNF or other conditions such as cancer, infection, or autoimmune disease.

The compounds described herein which selectively inhibit the activity of the p38 α isoform are useful for treating conditions associated with activation of p38 α , in particular those associated with cardiac hypertrophy, ischemia or other environmental stress such as oxidation injury, hyperosmolarity or other agents or factors that activate p38 α kinase, or cardiac failure, for example, congestive heart failure, cardiomyopathy and myocarditis.


Compounds which exhibit this activity are of the formula




or



or



or



wherein R^1 , R^2 , R^3 , Z^1 , and Z^2 are as defined in claim 1.

The manner of administration and formulation of the compounds described herein will depend on the nature of the condition, the severity of the condition, the particular subject to be treated, and the judgement of the practitioner; formulation will depend on mode of administration. As these compounds are small molecules, they are conveniently administered by oral administration by compounding them with suitable pharmaceutical excipients so as to provide tablets, capsules, syrups, and the like. Suitable formulations for oral administration may also include minor components such as buffers, flavoring agents and the like. Typically, the amount of active ingredient in the formulations will be in the range of 5%-95% of the total formulation, but wide variation is permitted depending on the carrier. Suitable carriers include sucrose, pectin, magnesium stearate, lactose, peanut oil, olive oil, water, and the like.

The compounds useful in the invention may also be administered through suppositories or other transmucosal vehicles. Typically, such formulations will include excipients that facilitate the passage of the compound through the mucosa such as pharmaceutically acceptable detergents.

The compounds may also be administered topically, for topical conditions such as psoriasis, or in formulation intended to penetrate the skin. These include lotions, creams, ointments and the like which can be formulated by known methods.

The compounds may also be administered by injection, including intravenous, intramuscular, subcutaneous or intraperitoneal injection. Typical formulations for such use are liquid formulations in isotonic vehicles such as Hank's solution or Ringer's solution.

Alternative formulations include nasal sprays, liposomal formulations, slow-release formulations, and the like, as are known in the art.

Any suitable formulation may be used. A compendium of art-known formulations is found in Remington's Pharmaceutical Sciences, latest edition, Mack Publishing Company, Easton, PA. Reference to this manual is routine in the art.

5 The dosages of the compounds of the invention will depend on a number of factors which will vary from patient to patient. However, it is believed that generally, the daily oral dosage will utilize 0.001-100 mg/kg total body weight, preferably from 0.01-50 mg/kg and more preferably about 0.01 mg/kg-10 mg/kg. The dose regimen will vary, however, depending on the conditions being treated and the judgment of the practitioner.


10 As implicated above, although the compounds of the invention may be used in humans, they are also available for veterinary use in treating animal subjects.

The following examples are intended to illustrate but not to limit the invention.

15 Examples 1-3 illustrate Reaction Scheme 1:

Example 1


Preparation of 4-BOC piperazinyl-benzimidazole-5-carboxamide



20 Benzimidazole-5-carboxylic acid (3.25g, 20 mMol) was reacted with 2.52g (20 mMol) diisopropylcarbodiimide in dry DMF at room temperature for 15 minutes. To this reaction mixture was added 3.75g (20 mMol) *t*-butyl-1-piperazine carboxylate, and the mixture was stirred for 18h. The mixture was poured into water and extracted with methylene chloride (3x100 mL). The combined extracts were washed again with water, brine and dried over MgSO₄. After removal of the solvent *in vacuo*, the residue
25 was chromatographed on a column of silica gel eluting with CHCl₃-Methanol (gradient, methanol 0 to 5 %) to yield 5.69g (86 %) of the product. ¹H-NMR (DMSO d₆): s 8.3 (1H); m 7.7-7.6 (2H), m 7.2-7.3 (1H), m 3.6-3.3 (8H) s 1.4 (9H); MS (ESI) *m/e* 330 (*m*⁺).

Example 2


Preparation of piperazinyl-benzimidazole-5-carboxamide



5 The N-BOC piperazinyl-benzimidazole-5-carboxamide (5.6 g) was stirred in 20 mL 4 Molar HCl-dioxane for 1 h. The dioxane was removed under reduced pressure to yield the hydrochloride salt in quantitative yield. This was used for alkylations without any further purifications.

Example 3

Preparation of 4-(2,6-difluorobenzyl)-piperazinyl-benzimidazole-5-carboxamide



10 A. Piperazinyl-benzimidazole-5-carboxamide (0.186g, 0.5 mMol) was taken in 5 mL DMF and 0.101g (1 mMol) triethylamine was added and stirred for 15
15 minutes at room temperature. To this reaction mixture was added 0.104 g 2,6-difluorobenzyl bromide and the mixture was stirred for 20 h. This was poured into water and extracted with methylene chloride (3x50 mL). The combined extract was further washed with brine, water and dried over MgSO_4 . The solvent was removed *in vacuo* and the residue chromatographed on silica gel eluting with chloroform-
20 methanol (0 to 5% methanol, gradient). Evaporation of the desired fraction gave 48.9 mg of the desired product; MS(ESI) *m/e* 356 (M^+).

 B. Using the procedure set forth in paragraph A, the following compounds were prepared:

Preparation of	By substituting for 2,6-difluorobenzyl bromide :
4-(2,3-difluorobenzyl)-piperazinyl-benzimidazole-5-carboxamide MS (ESI) m/e 356 (M^+)	2,3-difluorobenzyl bromide
4-(3,5-difluorobenzyl)-piperazinyl-benzimidazole-5-carboxamide MS (ESI) m/e 356 (M^+)	3,5-difluorobenzyl bromide
4-(3-chlorobenzyl)-piperazinyl-benzimidazole-5-carboxamide MS (ESI) m/e 354 (M^+)	3-chlorobenzyl bromide
4-(4-carboxymethyl benzyl)-piperazinyl-benzimidazole-5-carboxamide MS (ESI) m/e 378 (M^+)	methyl-4-(bromomethyl)-benzoate
4-(4-methoxybenzyl)-piperazinyl-benzimidazole-5-carboxamide MS (ESI) m/e 350 (M^+)	4-methoxybenzyl chloride
4-(4-trifluoromethoxybenzyl)-piperazinyl-benzimidazole-5-carboxamide MS (ESI) m/e 404 (M^+)	4-(4-trifluoromethoxy)-benzyl bromide
4-(4-methylbenzyl)-piperazinyl-benzimidazole-5-carboxamide MS (ESI) m/e 334 (M^+)	4-methylbenzyl bromide
4-(2,4-dichlorobenzoyl)-piperazinyl-benzimidazole-5-carboxamide MS (ESI) m/e 403 (M^+)	2,4-dichlorobenzoyl chloride
4-(3,4-dichlorobenzoyl)-piperazinyl-benzimidazole-5-carboxamide MS (ESI) m/e 403 (M^+)	3,4-dichlorobenzoyl chloride
4-[trans-3-(trifluoromethyl)-cinnamoyl]-piperazinyl-benzimidazole-5-carboxamide MS (ESI) m/e 428 (M^+)	trans-3-(trifluoromethyl)-cinnamoyl chloride
4-(4-chlorobenzoyl)-piperazinyl-benzimidazole-5-carboxamide MS (ESI) m/e 369 (M^+)	4-chlorobenzoyl chloride
4-benzoylpiperazine-benzimidazole-5-carboxamide MS (ESI) m/e 334 (M^+)	benzoyl chloride
4-(2-trifluoromethylbenzoyl)-piperazinyl-benzimidazole-5-carboxamide MS (ESI) m/e 402 (M^+)	2-(trifluoromethyl)-benzoyl chloride
4-(4-methoxybenzoyl)-piperazinyl-benzimidazole-5-carboxamide MS (ESI) m/e 364 (M^+)	4-methoxybenzoyl chloride

Example 4 illustrates Reaction Scheme 2:

Example 4

Preparation of 4-(3,4-dichlorophenyl)-piperazinyl-benzimidazole-5-carboxamide

- 5 A. Benzimidazole-5-carboxylic acid (1 mMol, 162 mg) was dissolved in 5 mL dry DMF and reacted with 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide hydrochloride for 15 minutes. 1-(3,4-dichlorophenyl)-piperazine, 1 mMol (231 mg) was added followed by 10 mg DMAP. The mixture was stirred for 20 h at room temperature. The reaction mixture was poured into water and extracted with

methylene chloride (3x 50 mL). The extracts were combined, washed with brine, water and dried over MgSO_4 . After evaporation of the solvent, the residue was chromatographed on silica gel with chloroform-methanol (0-5 % methanol, gradient). Evaporation of the desired fractions gave 150 mg (40 %) of the title compound; MS (ESI) m/e 375 (M^+).


B. Using the procedure of paragraph A, the following were prepared:

Preparation of	Substituting for 1-(3,4-dichlorophenyl) piperazine,
4-(4-chlorobenzhydryl)-piperaziny-benzimidazole-5-carboxamide MS (ESI) m/e 431 (M^+)	1-(4-chlorobenzhydryl)-piperazine
4-trans-1-cinnamyl piperaziny-benzimidazole-5-carboxamide MS (ESI) m/e 346 (M^+)	trans-1-cinnamyl piperazine
4-[bis(4-fluorophenyl)-methyl]-piperaziny-benzimidazole-5-carboxamide MS (ESI) m/e 432 (M^+)	1-Bis (4-fluorophenyl)-methyl piperazine
4-(4-chlorobenzyl)-piperaziny-benzimidazole-5-carboxamide MS (ESI) m/e 354 (M^+)	1-(4-chlorobenzyl)-piperazine
4-(2-chlorobenzyl)-piperaziny-benzimidazole-5-carboxamide MS (ESI) m/e 354 (M^+)	1-(2-chlorobenzyl)-piperazine
4-benzylpiperaziny-benzimidazole-5-carboxamide MS (ESI) m/e 320 (M^+)	1-benzyl piperazine

Example 5 illustrates Reaction Scheme 3:

Example 5

10 A. Preparation of 4-(4-methylthiobenzyl)-piperaziny-benzimidazole-5-carboxamide



A mixture of 4-(methylthio)-benzaldehyde, 305 mg (2 mMol) and N-BOC piperazine, 372 mg (2 mMol) was stirred in dry methanol for 30 minutes. To this mixture was added 1.6 g of polymer-supported borohydride (2.5 mMol/g, on
15 Amberlite, IRA-400, Aldrich) and the mixture was stirred for 24 h. The polymer was removed by filtration and evaporation of the solvent yielded the 4-BOC- 1-(4-methylthio)-benzylpiperazine in quantitative yield. MS (ESI) m/e 322, (M^+).

The 4-BOC-1-(4-methylthio)-benzylpiperazine was taken in 10 mL 1:1 TFA/methylene chloride and stirred for 1 h at room temperature. The solvents were removed *in vacuo* and the residue was used without purification for coupling with benzimidazole-5-carboxylic acid.


- 5 Benzimidazole-5-carboxylic acid (2 mMol, 324 mg) was taken in 15 mL dry DMF and reacted with 2 mMol (382 mg) EDAC at room temperature for 15 minutes. The above described 1-(4-methylthio)-benzylpiperazine was added as a DMF solution followed by 505 mg (5 mMol) TEA. The mixture was stirred for 20 h. The mixture was poured into water and extracted with methylene chloride (3x 50 mL). The combined extracts were washed with brine, water and dried over MgSO₄. The solvent was removed *in vacuo* and the residue was chromatographed. Evaporation of the desired fractions gave the title compound; MS (ESI) *m/e* 366 (M⁺).
- 10

Using this procedure, the following were prepared:

Preparation of	Substituting for 4-(methylthio)-benzaldehyde
4-(3,4,5-trimethoxybenzyl)-piperazynyl-benzimidazole-5-carboxamide MS (ESI) <i>m/e</i> 410 (M ⁺)	3,4,5-methoxybenzaldehyde
4-(4-diethylaminobenzyl)-piperazynyl-benzimidazole-5-carboxamide MS (ESI) <i>m/e</i> 391 (M ⁺)	4-diethylaminobenzaldehyde
4-(biphenylmethyl)-piperazynyl-benzimidazole-5-carboxamide MS (ESI) <i>m/e</i> 396 (M ⁺)	4-biphenylcarboxaldehyde
4-(4-Phenoxybenzyl)-piperazynyl-benzimidazole-5-carboxamide MS (ESI) <i>m/e</i> 412 (M ⁺)	4-phenoxybenzaldehyde

15

B. Preparation of 4-benzyl-piperidinyl-benzimidazole-5-carboxamide



- Benzimidazole-5-carboxylic acid (1.62g, 10 mMol) was reacted with EDAC (1.92g, 10 mMol) in 40 mL dry DMF at room temperature for 15 minutes. To the reaction mixture was added 4-benzylpiperidine (1.75g, 10 mMol) and DMAP (~20mg, catalyst) and the mixture was stirred at room temperature for 20 h. It was
- 20


poured into water and extracted with methylene chloride (3x100 mL). The combined extract was washed with water, brine and again with water. The extract was dried over MgSO_4 and evaporated. The residue was chromatographed on a column of silica gel with chloroform-methanol (0 to 5 % methanol). Evaporation of the desired fractions gave 1.5g (47 %) of the product after recrystallization from ethyl acetate-hexane. ^1H NMR (CDCl_3): δ = 7.8 (s, 1H); 7.1-7.3 (m, 8H); 4.8-4.7 (broad m, 1H), 3.7-3.9 (broad m, 1H); 3.1-2.7 (broad m, 2H); 2.55 (d, 2H); 2.0-1.1 (m, 5H). MS (ESI) m/e 319 (M^+), 318 ($\text{M}^+ - \text{H}$).

10

Example 6

Preparation of Additional Benzimidazole Piperidinyl Embodiments

The reaction scheme in this example is generally as follows:



15

- Nitric acid/Sulfuric acid, 100°C, 1 hr.
- Methanol, 10% PdC, Formic acid, 1-3 hr.
- 90% aq. formic acid, reflux, 1.5 hr.
- Benzylpiperidine, EADC•HCl, DMAP, DMF.

2-methoxy-3,4-dinitrobenzoic acid:

20

4-Nitro-2-methoxybenzoic acid, 3.09 g, was added to 20 mL Nitric acid:Sulfuric acid 1:1, at 0°C. After addition was complete the reaction mixture was heated at 100 °C for 30 minutes. Cooled to room temperature and poured into 200 mL ice water. The aqueous layer was extracted with ethyl acetate and washed with saturated sodium chloride, dried over anhydrous sodium sulfate and concentrated to give a yellow solid. This material was purified by chromatography on silica using ethyl acetate/hexane/methanol/acetic acid 5/5/1/0.1. The yellow solid obtained was

25

used for the next step. EIMS M^+ 242 (Exp. 242). NMR, d_6 DMSO: s (1H) 8.5, s (1H) 7.95, s (3H) 4.05


2-methoxy-3,4-diaminobenzoic acid:

5 2-methoxy-3,4-dinitrobenzoic acid (1.0 g) was dissolved in methanol (50 mL) and treated with 100 mg 10% palladium on carbon. The reaction mixture was purged with nitrogen and placed in an ice bath. Upon treatment with 5 mL formic acid, brisk effervescence was noticed which subsides upon further cooling. The reaction mixture was filtered through celite and concentrated to give a tan solid. (Decolorization occurs
10 rapidly upon keeping) EIMS M^+ 182, Exp. 182.


6-methoxy-5-benzimidazole carboxylic acid:

 2-methoxy-3,4-diaminobenzoic acid, (0.5 g) was dissolved in 10 mL 90% aqueous formic acid. The mixture was brought to reflux and maintained there for 90
15 mins. Cooled to room temperature and the solvent removed under pressure to give a dark solid. EIMS M^+ 192, Exp. 192.

6-methoxy-(4-benzylpiperidiny)benzimidazole-5-carboxamide:




20 6-methoxy-5-benzimidazole carboxylic acid, (1 equivalent) was treated with 1.1 equivalent of EDAC·HCl and 1 equivalent of 4-benzylpiperidine in the presence of a catalytic amount of DMAP in DMF/DCM 1:1 for 3-6 hrs. The reaction mixture was then concentrated and taken up in ethyl acetate. After washing with 5% aq. Sodium carbonate and a solution of saturated sodium chloride, the organic layer was
25 dried over anhydrous sodium sulfate and concentrated to give crude material. This crude material was purified by chromatography on silica. M^+ 349, Exp. 349.

6-chloro-(4-benzylpiperidinyl)benzimidazole-5-carboxamide:


was prepared similarly. MH^+ 353, Exp. 353.

5

Example 7N-propylation of 4-benzyl-piperidinyl-benzimidazole-5-carboxamide

4-(4-Benzyl)-piperidinyl-benzimidazole-5-carboxamide (318 mg, 1 mMol) was taken in 20 mL acetone. KOH (solid, 280 mg, 5 mMol) was added followed by 2-iodopropane (1g ~ 6 mMol) and the mixture was refluxed for 20 h. The acetone was removed *in vacuo* and the residue extracted from water with methylene chloride (3x50 mL). The extract was dried, evaporated and the residue chromatographed on silica gel with $CHCl_3$ -Methanol (0 to 3 % methanol). MS (ESI) m/e 360 (M^+). HPLC: (Vydac C18 column, 5 to 40 % acetonitrile/water containing 0.1 % TFA) two peaks showing both isomers.


15

Example 8Preparation of 4-benzylpiperidinyl-indole-5-carboxamide.


Indole-5-carboxylic acid (1.61g, 10 mMol) was reacted with EDAC (1.92g, 10 mMol) in 40 mL dry DMF for 15 minutes. 4-Benzylpiperidine (1.75g, 10 mMol) was added followed by DMAP (20 mg, catalyst) and the reaction mixture was stirred for 20 h. The mixture was poured into water and extracted with methylene chloride (3x100 mL). The combined extract was washed with dilute hydrochloric acid, saturated sodium bicarbonate and water and dried over MgSO₄. After evaporation of the solvent, the residue was chromatographed with methylene chloride-methanol (0 to 2 % methanol, gradient) to yield 1.60 g (50 %) of the product after recrystallization from ether-Hexane. MS (ESI) *m/e* 318 (M⁺), (317⁺-H). ¹HNMR (CDCl₃) δ=8.5 (s, 1H); 7.7 (s, 1H); 7.4-7.15 (m, 8H); 6.8 (s, 1H); 4.8-4.6 (br, m, 1H); 4.1-3.9 9br, m, 1H); 3.1-2.7 (br, m, 2H); 2.6 (d, 2H); 1.9-1.7 (br, m, 3H); 1.4-1.2 (br, m, 2H).

Example 9

Preparation of 4-benzylpiperidinyl-1-(2-propyl)-indole-5-carboxamide



A mixture of 4-benzylpiperidinyl-benzimidazole-5-carboxamide (318 mg, 1 mMol), solid KOH (280 mg, 5 mMol) and 2-iodopropane (1g, 6 mMol) was refluxed in 20 mL acetone for 20 h. After the removal of acetone *in vacuo*, the residue was extracted from water with methylene chloride (3x50 mL). The combined extract was dried, evaporated and chromatographed to yield 180 mg (50 %) of the desired product. ¹HNMR (CDCl₃): δ=7.7 (s, 1H); 7.4-7.1 (m, 7H); 4.8-4.6 (m, 1H); 3.0-2.7 (br, m, 4H); 2.6 (d, 2H); 1.8-1.45 (m, 3H); 1.5 (d, 6H); 1.3-1.1 (m, 2H). MS (ESI) *m/e* 360 (M).

Example 10Preparation of 4-(4-chlorobenzyl)-piperazinyl-1-(2-propyl)-indole-5-carboxamide

4-(4-Chlorobenzyl)piperazinyl-indole-5-carboxamide (420 mg, 1.32 mMol)
 5 was taken in acetone. Solid KOH (280 mg, 5 mMol) was added followed by the
 addition of 2-iodopropane (1g, 6 mMol) and the mixture was refluxed with stirring for
 20 h. Acetone was removed *in vacuo* and the residue was extracted from water using
 methylene chloride. The extract was dried and evaporated and the residue was
 chromatographed on a column of silica gel using ethylacetate-hexane (ethylacetate 0
 10 to 25 %, gradient) and recrystallized from ether-hexane to yield 300 mg of the
 product. ¹HNMR (CDCl₃): δ= 7.6 (s, 1H); 7.3-7.1 (m, 6H); 6.5 (s, 1H); 4.65-4.55 (m,
 1H); 3.8-3.5 (m, 4H); 3.4 (s, 2H); 2.4-2.5 (s, 4H); 1.5 (d, 6H). MS (ESI) *m/e* 395
 (M⁺).

15

Example 11Preparation of Additional Analogs

Using the procedure of Example 8, following are prepared:

Preparation of	Using the procedure set forth in Example 8, but substituting for indole-5-carboxylic acid
4-benzylpiperidinyl-indole-6-carboxamide MS (ESI) <i>m/e</i> 318 (M ⁺), (317 ⁺ -H)	Indole-6-carboxylic acid
4-benzylpiperidinyl-benzotriazole-5-carboxamide MS (ESI) <i>m/e</i> 320 (M ⁺), (319 ⁺ -H)	Benzotriazole-5-carboxylic acid

Using the procedure of Example 10, these compounds were alkylated, e.g.,

Preparation of	Using the procedure set forth in Example 11, but substituting for 4-chlorobenzyl piperidinyl indole-5-carboxamide
4-benzylpiperidinyl-1-(2-propyl)-indole-6-carboxamide MS (ESI) <i>m/e</i> 360 (M^+)	4-benzyl piperidinyl-indole-6-carboxamide
4-benzylpiperidinyl-1-(2-propyl)-benzotriazole-5-carboxamide	4-benzyl piperidinyl-benzotriazole-5-carboxamide

Example 12

Preparation of 3-Chlorobenzylpiperazinyl-N-benzyl-benzimidazole-5- and 6-carboxamides

5

A. This paragraph describes the procedure for formation of the N-benzyl derivatives of the compounds of the invention; succeeding paragraphs describe alkylation with other moieties.

3-Chlorobenzylpiperazinyl-benzimidazole-5-carboxamide (0.12 g, 0.33 mMol) and the benzyl bromide (0.058 g, 0.33 mMol) in 15 mL DMF were combined with K_2CO_3 (0.09 g, 0.66 mMol). The mixture was stirred at RT overnight, then heated at 45°C for 3 h. EtOAc was added and washed with water. The organic layer was evaporated and the isomers were separated by silica gel column chromatography using 5% MeOH in EtOAc. of isomer a (70 mg, 48%), MS (ESI) *m/e* 444 (M^+) and of isomer b (40 mg, 27 %), MS (ESI) *m/e* 444 (M^+) were obtained.

15

Similar treatment of the 6-carboxamide yields the corresponding compound where R^2 is benzyl.

B. 3-Chlorobenzylpiperazinyl-N-(2-propyl)-benzimidazole-5- and 6-carboxamides.

20

3-Chlorobenzylpiperazinyl-benzimidazole-5-carboxamide was alkylated substituting 2-iodopropane for benzyl bromide in paragraph A. The isomers were separated using the same chromatographic conditions. Isomer a, MS (ESI) *m/e* 396 (M^+); isomer b, MS (ESI) *m/e* 396 (M^+).

Similar treatment of the 6-carboxamide yields the corresponding compound where R^2 is 2-propyl.

25

C. 3-Chlorobenzylpiperazinyl-N-methyl-benzimidazole-5- and 6-carboxamide

3-Chlorobenzylpiperazinyl-benzimidazole-5-carboxamide was alkylated substituting iodomethane for benzyl bromide in the procedure of paragraph A. The
5 isomers were separated using silica gel column chromatography using 50% acetone in acetonitrile as the eluting solvent. Isomer a, MS (ESI) m/e 368 (M^+), isomer b, MS (ESI) m/e 368 (M^+).

Similar treatment of the 6-carboxamide yields the corresponding compound where R^2 is methyl.

10 Similarly, 4-benzylpiperidinyl-(1-methyl)-indole-5-carboxamide (MS (ESI) m/e 332 (M^+)) was prepared from 4-benzylpiperidinyl-indole-5-carboxamide.

Similar treatment of the 6-carboxamide yields the corresponding compound where R^2 is methyl.

15 D. 3-Chlorobenzylpiperazinyl-N-ethyl-benzimidazole-5- and 6-carboxamides


3-Chlorobenzylpiperazinyl-benzimidazole-5-carboxamide was alkylated substituting iodoethane for benzyl bromide in paragraph A. Isomer a, MS (ESI) m/e 382 (M^+); isomer b, MS (ESI) m/e 382 (M^+).

20 Similar treatment of the 6-carboxamide yields the corresponding compound where R^2 is ethyl.

Similarly, 4-benzylpiperidinyl-(1-ethyl)-indole-5-carboxamide (MS (ESI) m/e 346 (M^+)) was prepared from 4-benzylpiperidinyl-indole-5-carboxamide.


Similar treatment of the 6-carboxamide yields the corresponding compound where R^2 is ethyl.

25

Example 13Preparation of 4-(4-chlorobenzyl)-piperidiny-indole-5-carboxamide

This example illustrates Reaction Scheme 5.

5 A. Preparation of N-BOC-4-(4-chlorobenzylene)-piperidine



N-BOC-4-piperidone (2.0 g; 10 mmol) was taken with diethyl-4-chlorobenzylphosphonate (2.6 g; 10 mmol) in dry THF. Sodium hydride (400 mg, 60 % dispersion in mineral oil; 10 mmol) was added and the mixture was refluxed for
10 three hr. The THF was removed in vacuo and the residue extracted from water with methylene chloride. The extract was dried over MgSO_4 , evaporated and the residue was chromatographed on silica gel to yield 0.615 g of the desired product. ^1H NMR (CDCl_3): $\delta=7.3$ (d, 2H); 7.1 (d, 2H); 6.3 (s, 1H); 3.55-3.50 (m, 2H); 3.45-3.35 (m, 2H); 2.45-2.35 (m, 2H); 2.30-2.25 (m, 2H); 1.25 (s, 9H). EIMS: 307 (M^+), 251 (M^+ -
15 C_3H_8).

B. Coupling of 4-Chlorobenzylene piperidine with indole-5-carboxylic acid

The N-BOC-4- (4-chlorobenzylene)-piperidine, described above, was deprotected by stirring in 20 mL 1:1 dichloromethane-trifluoroacetic acid for 1 h. It
20 was evaporated and dried in vacuo for 1 h to remove all traces of trifluoroacetic acid. It was redissolved in 15 mL dichloromethane and the TFA salt was neutralized by the addition of a slight excess of triethylamine. Solution A.

Indole-5-carboxylic acid 0.32g (2 mmol) was reacted with 0.383g EDAC in 30 mL dry dichloromethane for 15 minutes. To this solution was added the methylene chloride solution of 4-chlorobenzyl-piperidine (solution A) followed by the addition of 10 mg of DMAP. The mixture was stirred for 20 h. The mixture was washed with water, 2N HCl, 5% sodium carbonate and then water. The organic solution was dried, evaporated and the residue was chromatographed on silica gel eluting with ethylacetate-hexane (1:4). Yield: 260 mg (37 %). EIMS: 350 (M^+), 315 (M^+-Cl) 1H NMR ($CDCl_3$): δ =8.4 (s, 1H); 7.7 (s, 1H); 7.3-7.0 (m, 7H); 6.5 (s, 1H); 6.25 (s, 1H); 3.8-3.0 (m, br, 4H); 2.6-2.20 (m, br, 4H).

C. Hydrogenation of 4-(4-chlorobenzyl)-piperidine-indole-5-carboxamide


4-(4-Chlorobenzyl)-piperidine-indole-5-carboxamide (240 mg, 0.68 mmol) was dissolved in 40 mL THF. Pd/C (25 mg) was added and the mixture was hydrogenated (1 atm) for 20 h with rapid stirring. The catalyst was removed by filtration through celite and the organic solution was evaporated and the residue was recrystallized from methylene chloride/hexane. Quantitative yield. EIMS: 352 (M^+), 351 (M^+-H).

Example 14

Using the general procedure set forth in Example 13, the following are prepared:

Preparation of	Substituting for 4-chlorobenzyl piperidine
4-(3-chlorobenzyl)-piperidiny-indole-5-carboxamide MS (ESI) m/e 353 (M^+)	3-chlorobenzyl piperidine
4-(2-chlorobenzyl)-piperidiny-indole-5-carboxamide MS (ESI) m/e 353 (M^+)	2-chlorobenzyl piperidine

Example 15

Synthesis of cis-2-Methyl-4-benzylpiperidin-1-yl-indole-5-carboxamide

- A. A mixture of 4-benzylpiperidine (3.52 mL, 20.0 mMol) and di-tert-butyl-dicarbonate (5.45 g, 25.0 mMol) in 100 mL of THF was refluxed for 20 h. After cooling to rt the reaction mixture was poured into water and extracted with ethyl acetate (2 x 100 mL). the combined organic extract was washed with water and brine. The extract was dried over Na₂SO₄ and evaporated. The residue was chromatographed on a column of silica gel with 10% ethyl acetate-hexane.
- 10 Evaporation of the desired fractions gave 5.02 g (91%) of the product as an oil. MS (ESI) *m/e* 275 (M⁺).
- B. A mixture of 1-BOC-4-benzylpiperidine (0.825 g, 3.0 mMol) and N,N,N',N'-tetramethylethylenediamine (TMEDA) (0.59 mL, 3.9 mMol) in 6 mL of Et₂O was cooled to -78°C under argon. A 1.3M solution of s-BuLi in cyclohexane
- 15 (3.0 mL, 3.9 mMol) was added dropwise. After the addition was complete, the reaction mixture was stirred at -20°C for 30 min and cooled back to -78°C. Methyl iodide (0.28 mL, 4.5 mMol) was added and the reaction mixture was stirred at -78°C for 5 min, the cooling bath removed and stirring was continued an additional 3 min. The reaction mixture was poured into water and extracted with ethyl acetate (2 x 25
- 20 mL). The combined organic extract was washed with water and brine. The extract was dried over Na₂SO₄ and evaporated to give 0.58 g (67%) of an oil that was one spot by TLC (silica gel, 10% ethyl acetate-hexane). This material was used directly in the next step. MS (ESI) *m/e* 289 (M⁺).
- C. To a solution of 1-BOC-2-methyl-4-benzylpiperidine (0.29 g, 1.0
- 25 mMol) in 5 mL of dichloromethane was added trifluoroacetic acid (TFA) (0.5 mL). After stirring at rt for 10 h the reaction mixture was evaporated *in vacuo* and


- 34 -

azcotroped twice with dichloromethane and twice with hexane. The residue was dissolved in 5 mL of dichloromethane and diisopropylethylamine (1.6 mL, 10 mMol) was added. In a separate flask a mixture of 5-indolecarboxylic acid (0.19 g, 1.2 mMol) and EDAC (0.23 g, 1.2 mMol) was dissolved in 15 mL of dichloromethane and stirred at rt for 5 min. To this reaction mixture was added the first solution, and the resulting mixture stirred at rt for 20 h. The reaction mixture was poured into water and extracted with ethyl acetate (2 x 50 mL). The combined organic extract was washed with water and brine. The extract was dried over Na_2SO_4 and evaporated. The residue was chromatographed on a column of silica gel with 1% MeOH-dichloromethane. Evaporation of the desired fractions gave 0.18 g (54%) of the product as an oil.


When tested as described below, the title compound has an IC_{50} =280 nM.

Example 16

Preparation of 4-Chloro-(4-benzylpiperidiny)indole-5-carboxamide and 6-Chloro-(4-benzylpiperidiny)indole-5-carboxamide



and



A. The indole synthesis was accomplished by the method of Gassman, P. G. *J Am Chem Soc* (1974) 96: 5495-5507. To a solution of 2.0 g (10.8 mmol) of 4-amino-2-chloro methylbenzoate in 30 mL CH_2Cl_2 at -60°C was added 1.2 g (10.8 mmol) of t-butyl hypochlorite in 20 mL of CH_2Cl_2 . After 10 min. 10.8 mmol of methylthioacetaldehyde dimethyl acetal in 10 ml of CH_2Cl_2 was added, stirring at -60°C continued for 1 h. Subsequently, 10.8 mmol of Et_3N in 10 ml CH_2Cl_2 was added and the solution was allowed to warm to room temperature. The solvent was evaporated and the residue was dissolved in 30 mL CCl_4 , 5 mL of Et_3N was added and the mixture was refluxed for 4 h. The solvent was removed and the residue was dissolved in 50 mL of ether. Cyclization of the acetal to indole was effected by

stirring this solution for 3 h with 20 mL of 2N HCl. The ethereal layer was washed with saturated NaHCO₃, dried, filtered and evaporated. The isomeric indoles were separated by column chromatography on silica gel. The structure of the isomers was identified by NMR spectroscopy. Isomer a: 5-carboxymethyl-4-chloro-3-thiomethylindole, ¹H NMR (CDCl₃) δ 2.35 (s, 3H), 3.95 (s, 3H), 7.32 (s, 1H), 7.42 (s, 1H), 8.33 (s, 1H), 8.61 (s, 1H). Isomer b: 5-carboxymethyl-6-chloro-3-thiomethylindole, ¹H NMR (CDCl₃) δ 2.42 (s, 3H), 3.97 (s, 3H), 7.20 (s, 1H), 7.25 (d, 1H), 7.71 (d, 1H), 8.58 (s, 1H).

A solution of 100 mg of 5-carboxymethyl-4-chloro-3-thiomethylindole (isomer a) in 10 mL of ethanol was treated with W-2 Raney nickel until dethiomethylation was completed. The indole ester isolated was treated with NaOH in methanol:water (1:1) and thereby 60 mg of 4-chloroindole-5-carboxylic acid was isolated as a white solid, ¹H NMR (DMSO-d₆) δ 6.61 (s, 1H), 7.41 (d, 1H), 7.52 (s, 1H), 7.62 (d, 1H), 11.62 (s, 1H).


To a solution of 50 mg (0.25 mmol) of above indole acid in 10 mL of DMF was added 50 mg (0.28 mmol) of 4-benzylpiperidine and 60 mg (0.28 mmol) of EDAC. The reaction mixture was stirred overnight, diluted with ethyl acetate and washed with water. the organic layer was dried, filtered and evaporated to get a white solid. This was purified by silica gel chromatography followed by crystallization to obtain 50 mg of 4-chloro-5-(4-benzylpiperidinyl)-indole carboxamide as a white solid, MS (M⁺ 352).

Isomer b was converted to 6-chloroindole-5-carboxylic acid using the same reaction sequence as described above and was coupled to 4-benzylpiperidine to obtain 6-chloro-5-(4-benzylpiperidinyl)-indole carboxamide as a white solid, MS (M⁺ 352).

B. Using the method of paragraph A, 4-Chloro-(4-(4-fluorobenzyl)piperidinyl)indole-5-carboxamide and 6-Chloro-(4-(4-fluorobenzyl)piperidinyl)indole-5-carboxamide were prepared.

Example 17The Corresponding 6-Piperidinyl Indole Derivatives.

Compounds similar to those in paragraphs A and B of Example 16, but wherein the piperidinyl substituent is at the 6-position are synthesized as follows:



5

a) Methanol, thionyl chloride, reflux

b)

i) N-chlorosuccinimide, DCM, methylthioacetaldehyde dimethylacetal, triethylamine, reflux

10

ii) CHCl_3 , reflux

iii) HCl

c) Raney Ni, EtOH

d) Methanol, sodium hydroxide, reflux


e) Benzylpiperidine, EDAC·HCl, DMAP, DMF/DCM.

15



f) Acetone, potassium hydroxide, nicotinoyl chloride.

Specifically, the following compounds were prepared according to this method:


- 37 -

4-benzylpiperidiny-5-chloroindole-6carboxamide:MH⁺ 351, exp 352;

5

4-benzylpiperidiny-7-chloroindole-6-carboxamide:MH⁺ 351, exp 352;1-nicotinoyl-4-benzylpiperidiny-7-chloroindole-6-carboxamide:

10

MH⁺ 457, exp 457;1-nicotinoyl-3-(2-dimethylamino)ethyl amino carbonyl-4-benzylpiperidiny-7-chloroindole-6-carboxamide:


15

MH⁺ 571, exp 571.


(See Examples 19-21 for addition of substituents at positions 1 and 3.)

Example 18

5 Preparation of 4-Methoxy-(4-benzylpiperidinyl)indole-5-carboxamide and 6-Methoxy-(4-benzylpiperidinyl)indole-5-carboxamide



10 A. Preparation of 4-Methoxyindole and 6-methoxyindole-5-carboxylic acids.




The corresponding methyl esters of these indole acids were prepared by a modified method of Scheme 6 according to Inoue, S. *Heterocycles*, (1992) 34: 1017-1029, wherein the two isomeric indole acids were obtained in a 3: 2 ratio. 5-carboxymethyl- 4-methoxyindole. ¹H NMR (CDCl₃) δ 2.42 (s, 3H), 3.92 (s, 3H), 4.13 (s, 3H), 7.14 (d, 1H), 7.18 (d, 1H), 7.55 (d, 1H), 9.41 (s, 1H). 5-Carboxymethyl 6-methoxyindole. ¹H NMR (CDCl₃) δ 2.38 (s, 3H), 3.81 (s, 3H), 3.95 (s, 3H), 6.83 (s, 1H), 7.21 (s, 1H), 8.22 (s, 1H), 8.50 (s, 1H).

20 B. Conversion to Title Compound

4-Methoxy or 6-methoxy-indole-5-carboxylic acid was coupled with 4-benzylpiperidine to obtain the title compounds, MS (M⁺ 349).

In addition, 4-methoxy-indole-5-carboxylic acid was coupled with 4-(4-fluorobenzyl)piperidine to obtain 4-methoxy-(4-(4-fluorobenzyl) piperidinyl)indole-5-carboxamide, MS (M⁺ 367) and 6-methoxy-indole-5-carboxylic acid was coupled with 4-(4-fluorobenzyl)piperidine to obtain 6-methoxy-(4-(4fluorobenzyl) piperidinyl)indole-5-carboxamide. MS (M⁺ 367).


Example 19Preparation of N-(3-cyclohexylmethylamino-2-hydroxypropyl)-4-benzylpiperidinyl-indole-5-carboxamide

5

A. The title compound was prepared according to scheme 7. To an ice-cold solution of 1.0 g (3.0 mmol) of 4-benzylpiperidinyl indole-5-carboxamide in acetone was added 15 mmol of powdered KOH followed by 3.0 mmol of epibromohydrin and the mixture was stirred for 30 min. The mixture was filtered and
10 the solution was evaporated. The residue was dissolved in ethyl acetate, washed with water, dried and evaporated. After purification by silica gel column chromatography 435 mg of epoxide was obtained. MS (M^+ 373).


To a solution of 200 mg (0.54 mmol) of the above indole epoxide in 5 mL of MeOH was added 121 mg (1.1 mmol) of cyclohexylmethylamine and the mixture was
15 refluxed for 1 hr. The crude product was purified on silica gel column. The amino compound was then converted into its HCl salt by treating with ethanolic HCl. MS (M^+ 487).

B. Following the procedure of paragraph A, but substituting piperazine for cyclohexylmethylamine, N-(3-N-methylpiperazinyl-2-hydroxypropyl)-4-benzylpiperidinyl indole-5-carboxamide:
20




was prepared. MS (M^+ 473);

substituting benzylamine for cyclohexylmethylamine N-(3-benzylamino-2-hydroxypropyl)-4-benzylpiperidinyl indole-5-carboxamide:




MS (M+ 481) was prepared;

- 5 substituting p-methoxybenzylamine for cyclohexylmethylamine, N-[3-{(4-methoxybenzyl)-amino}-2-hydroxypropyl]-5-(4-benzylpiperidinyl indole-5-carboxamide:




MS (M+ 511) was prepared; and

- 10 substituting propylamine for cyclohexylmethylamine, N-{3-n-propylamino-2-hydroxypropyl}-4-benzylpiperidinyl indole-5-carboxamide:




MS (M⁺) 433 was prepared.

Example 20Preparation of Additional 1-Substituted DerivativesA. Preparation of N-(4-pyridoyl)-4-benzylpiperidiny indole-5-carboxamide


5 0.318 mg (1 mmol) of 4-benzylpiperidiny indole-5-carboxamide was dissolved in 15 mL dry DMF. 80 mg (60% suspension in oil) of sodium hydride was added and the mixture was stirred for 30 minutes under nitrogen. The mixture was cooled to 0° C and 200 mg (1.1 mmol) isonicotinyl chloride hydrochloride was added and the mixture was stirred for 20 hr at room temperature. The reaction was quenched
10 by the addition of sat. ammonium chloride solution, diluted with water and the product was extracted with dichloromethane. The extract was dried, evaporated and the residue was chromatographed on silica gel (ethyl acetate-hexane, gradient, 50-75 % ethyl acetate) to yield 150 mg of the pure product. ESI MS (M^+ 423, M^+-H , 422).

 Using the procedure of the previous paragraph, but substituting 4-picolyl
15 chloride hydrochloride for isonicotinyl chloride hydrochloride, N-(4-pyridylmethyl)-4-benzylpiperidiny indole-5-carboxamide:




MS (M^+ 409) was prepared.

 B. Preparation of 1-Nicotinoyl-(4-benzylpiperidiny)-indole-6-
20 carboxamide:




This compound was prepared similarly. M^+ 423.

C. Preparation of 1-nicotinoyl-6-methoxy-(4-benzylpiperidinyl)-indole-5-carboxamide:



5 This compound was prepared similarly. M^+ 490.

D. Preparation of N-methylacetyl-4-Benzylpiperidinyl indole-5-carboxamide and its free acid:



1.95 g (6.13 mmol) of 4-benzylpiperidinyl indole-5-carboxamide was
10 dissolved in 30 mL dry DMF and was treated with 320 mg (8 mmol, 60% suspension
in oil) sodium hydride for 30 minutes. The reaction mixture was cooled to 0° C and
1.225g (8 mmol) bromomethylacetate was added and stirring continued for 1 h at 0°
C. The ice-bath was removed and stirring continued for another 6 h at room
temperature. The reaction was quenched by the addition of sat. ammonium chloride
15 solution, diluted with water and extracted with dichloromethane. The extract was
dried, evaporated and the residue purified by column chromatography on silica gel
eluting with ethylacetate-hexane (25 -35% ethylacetate) to yield 2.2 g (92%) of the
desired product. MS: M^+ , 390; M^+-1 , 389. ^1H NMR (CDCl_3): δ 7.7 (s, 1H); 7.35-7.1
(m, 8H); 6.6 (s, 1H); 5.1 (s, 2H); 3.75 (s, 3H); 3.0-2.7 (br, m, 4H); 2.6 (d, 2H); 1.9-1.2
20 9m, 5H).


2.15 g, (5.5 mmol) of 4-Benzylpiperidinyl indole-5-carboxamide-1-
methylacetate from the previous paragraph was taken in 20 mL ethanol . A solution of
2.0 g K_2CO_3 in 20 mL water was added and the mixture was refluxed for 2h. The
ethanol was removed under reduced pressure, the remaining solution diluted with

water and acidified with conc. HCl. The precipitated product was collected by filtration, washed with water and dried to yield 1.9 g of the product. MS: M^+ , 376; $M^+ - H$, 375.

E. Preparation of 1-Acryloyl- (4-benzylpiperidiny)-indole-5-carboxamide:

5 0.318g (1 mmol) of (4-benzylpiperidiny)-indole-5-carboxamide was taken in 15 mL dry acetone and was reacted with 0.2g (5 mmol) of powdered KOH for 15 Min. The mixture was cooled in ice and 0.225 mg (2.5 mmol) of acryloyl chloride was added in one lot. Stirring continued at 0° C for 20 Min., after which the reaction was further stirred at room temp. for 1 h. the solvent was removed in vacuo and the
10 residue was extracted with ethyl acetate from water. The extract was dried and evaporated. TLC (ethyl acetate-hexane) and mass spectrum (M^+ at 372) confirmed the desired product. This product was used without further purification for the next step.


F. 1-[3-(2-propylamino)-propionyl]-(4-benzylpiperidiny)-5-indole carboxamide:



15


The above obtained product was dissolved in 20 mL dichloromethane and reacted with 0.1 mL isopropylamine at room temperature for 18 h. The solvent was removed and the product was purified by column chromatography on silica gel eluting with chloroform-methanol (95:5). Yield: 180 mg, M^+ , 431.

20 G. 1-(3-piperazinylpropionyl)-(4-benzyl)-piperidiny-indole-5-carboxamide:




The 1-acryloyl-(4-benzyl)-piperidiny-indole-5-carboxamide above was reacted with tert-butyl-1-piperazine carboxylate as described before. The product was deprotected using methanolic HCl. M^+ 458.

H. 1-(3-benzylaminopropionyl)-(4benzylpiperidiny)-indole-5-
5 carboxamide:




was prepared by reacting 1-acryloyl-(4-benzylpiperidiny)-indole-5-carboxamide with benzylamine. M^+ 479.

I. 1-(3-morpholinylpropionyl)-4-(4-benzylpiperidiny)-indole-5-
10 carboxamide



was prepared by reacting 1-acryloyl-(4-benzylpiperidiny)-indole-5-carboxamide with morpholine, M^+ 459.


J. Preparation of 4-benzylpiperidiny indole-5-carboxamide-1-acetic acid-
15 n-propylamide:



4-Benzylpiperidiny-indole-5-carboxamide-1-acetic acid (200 mg, 0.53 mmol) from Example 21 was reacted with 120 mg (0.61 mmol) EDC in 10 mL dry


dichloromethane for 30 minutes. n-propylamine (100 μ L, excess) was added and the mixture stirred for 20 h. The solution was diluted with dichloromethane, washed with water and 5% sodium carbonate solution. The organic solution was dried and evaporated and the residue purified by silica gel chromatography with ethyl acetate-hexane (3:2) to yield 100 mg of the product. MS (M^+ 417).

K. Preparation of 4-benzylpiperidiny l-indole-5-carboxamide-1-acetic acid, (4-methoxybenzyl) amide:



Following the procedure in the previous paragraph but substituting p-methoxybenzylamine for n-propylamine, 4-benzylpiperidiny l-indole-5-carboxamide-1-acetic acid, (4-methoxybenzyl) amide was prepared. MS, ESI: $M+H$, 496.


L. Preparation of 1-(Diethylaminoethyl)-6-methoxy-(4-benzylpiperidiny l)-indole-5-carboxamide:



0.3g (0.862 mmol) of 6-methoxy-(4-benzylpiperidiny l)-indole-5-carboxamide was dissolved in 20 mL dry DMF. It was cooled in an ice-bath and reacted with 0.12 g NaH (3 mmol, 60 % suspension). 0.172 mg (1 mmol) of 2-(diethylamino)-ethylchloride hydrochloride was added and the mixture was stirred for 18 h. The reaction mixture was poured in to water and extracted with dichloromethane (3 x 75 mL). The combined extract was washed again with water, dried over anhydrous $MgSO_4$, evaporated and purified by silica gel chromatography on a chromatotron using CH_2Cl_2 -Methanol (95:5) to yield 0.22 g of the desired product. It was converted to the HCl salt and lyophilized, M^+ 448.


M. Following the procedure of the previous paragraph, but substituting 1-(diethylamino)-n-propylchloride hydrochloride for 1-(diethylamino)-ethylchloride

hydrochloride, 1-(Diethylamino) n-propyl- (4-benzylpiperidiny)- indole-5-carboxamide



was prepared. M^+ 468.


5 Similarly, 1-(Diethylamino)-ethyl-(4-benzylpiperidiny)-indole-5-carboxamide,



was prepared, M^+ 454.


Similarly, 1-(Diethylamino)-n-propyl-6-chloro-(4-benzylpiperidiny)-indole-5-carboxamide,

10



was prepared, M^+ 502.

Similarly, 1-(Diethylamino)-ethyl-(4'-fluoro-4-benzylpiperidiny)-indole-5-carboxamide,




15

was prepared, M^+ , 472.

Also 1-(Diethylamino)-n-propyl-6-methoxy-(4-benzylpiperidiny)-indole-5-carboxamide,

- 47 -




was prepared, M^+ , 498.

Example 21

Preparation of 3-Substituted Indoles

The general procedure for synthesis of the 3-substituted indoles is outlined as follows:




a) Trifluoroacetic anhydride, THF, 0°C, 3 hr.

b) aq. NaOH, reflux, 3-6 hrs.

c) R_2NH , EDAC·HCl, DMAP, DCM/DMF, 3-6 hrs.

A. 3-trifluoroacetyl-4-benzylpiperidinyl-indole-5-carboxamide: 4-benzyl

piperidinyl-indole-5-carboxamide (1 eq.) was dissolved in anhydrous THF. The reaction vessel was purged with nitrogen and placed in an ice bath. Trifluoroacetic anhydride (1.2-1.3 eq.) was added via syringe. The reaction was allowed to continue at 0°C, until no more starting material was discovered by thin layer chromatography. In some cases the addition of additional trifluoroacetic anhydride was required to facilitate the completion of the reaction. After completion of the reaction, the reaction mixture was concentrated and redissolved in the minimal amount of ethyl acetate for chromatography using silica. The crude material was chromatographed using ethyl acetate and hexanes (1:1). The identity of the product 3-trifluoroacetyl-4-benzyl piperidinyl indole-5-carboxamide was determined by electron impact mass spectroscopy. (MH^+ 413 (exp. 414), base peak 240.)













D. Similarly, making the appropriate substitutions for the carboxamide and for the reactive amine, the following compounds of the invention were prepared according to the reaction scheme set forth at the beginning of this example; all are of the formula set forth above, but with alternative substituents on the carbonyl moiety at the 3-position of the indole moiety as noted.

- i. 3-(2-methoxyethylaminocarboxamidyl)-(4-benzylpiperidinyl)indole-5-carboxamide (3-carbonyl substituent is 2-methoxy ethylamino, MH^+ 418, exp 418);
- ii. 3-(2-methylaminoethylaminocarboxamidyl)-(4-benzylpiperidinyl)indole-5-carboxamide (3-carbonyl substituent is 2-methylamino ethylamino, MH^+ 418, exp 418);
- iii. 3-(N-methyl-2-aminoethylaminocarboxamidyl)-(4-benzylpiperidinyl)indole-5-carboxamide (3-carbonyl substituent is 2-aminoethyl (methyl)amino, MH^+ 418, exp 418);
- iv. 3-(4-benzylpiperidinylcarboxamidyl)-(4-benzylpiperidinyl)indole-5-carboxamide (3-carbonyl substituent is 4-benzyl piperidinyl, MH^+ 519, exp 519);
- v. 3-(4-benzylpiperidinylcarboxamidyl)-(4-benzylpiperidinyl)indole-6-carboxamide (3-carbonyl substituent is 4-benzyl piperidinyl, MH^+ 519, exp 519);
- vi. 3-(4-fluorobenzylaminocarboxamidyl)-(4-benzylpiperidinyl)indole-5-carboxamide (3-carbonyl substituent is 4-fluorobenzylamino, MH^+ 469, exp 469);
- vii. 3-2-(3,4-dimethoxyphenyl)ethylaminocarboxamidyl-(4-benzylpiperidinyl)indole-5-carboxamide (3-carbonyl substituent is 2-(3,4-dimethoxyphenyl)ethylamino, MH^+ 525, exp 525);
- viii. 3-trifluoroacetyl-(4-benzylpiperidinyl)indole-5-carboxamide (3-carbonyl substituent is trifluoromethyl, MH^+ 413, exp 414);
- ix. 3-trifluoroacetyl-(4-benzylpiperidinyl)indole-6-carboxamide (3-carbonyl substituent is trifluoromethyl, MH^+ 413, exp 414);


x. 6-methoxy-3-(2-dimethylaminoethylamino)carboxamidyl-(4-benzylpiperidinyl)indole-5-carboxamide (3-carbonyl substituent is 2-dimethylaminoethyl, including also a 6-methoxy substituent in the 6-position, MH⁺ 462, exp 462).

5 The formulas of compounds i-x set forth above are shown below.




Alternatively, compounds of the invention which are derivatives of indole with substitutions at position 3 can be prepared using Reaction Scheme 8 set forth previously.

5 E. Preparation of 3-Morpholinomethyl-(4-benzylpiperidiny)-indole-5-carboxamide:



Using Scheme 8, 0.318 g (1mmol) of 4-benzylpiperidiny- indole-5-carboxamide, 0.1 g paraformaldehyde (3.3 mmol) and 0.1 mL morpholine was taken
10 in 25 mL abs. Ethanol and was acidified by the addition of 1 mL ethanolic HCl. The mixture was refluxed for 18h. The solvent was removed and the residue was extracted from 5% sodiumcarbonate solution with dichloromethane. The extract was dried, evaporated and the residue was purified by column chromatography using ethyl acetate-methanol (95:5), to yield 0.15 g of the desired product. It was converted to the
15 HCl salt and lyophilized. M^+ 454.

F. Preparation of Diethylaminomethyl-(4-benzylpiperidiny)-indole-5-carboxamide:



Using Scheme 8, this compound was prepared as follows. To a suspension of
20 0.41 g (1.28 mmol) 4-benzylpiperidiny-indole-5-carboxamide in 5 mL glacial acetic acid was added an ice cooled mixture of 1.2 mL aqueous formaldehyde (37%) and 0.16 mL of ethylamine (1.5 mmol). The reaction mixture was stirred for 30 Min. at 0° C and then continued stirring at RT for 18h. It was poured in to water, made basic by the addition of 20% sodium hydroxide solution and extracted with ethyl acetate. The

extract was dried and evaporated. The residue was purified by chromatography on silica gel eluting with chloroform-methanol-triethylamine (95:5:0.5) to yield 0.22 g of the title compound. MS: 403, M^+ ; 331, $M^+ - NEt_2$.

5

Example 22

Assay for p38 Kinase Inhibition

The compounds to be tested were solubilized in DMSO and diluted into water to the desired concentrations. The p38 kinase was diluted to 10 μ g/ml into a buffer containing 20 mM MOPS, pH 7.0, 25 mM beta-glycerol phosphate, 2 mg/ml gelatin,
10 0.5 mM EGTA, and 4 mM DTT.

The reaction was carried out by mixing 20 μ l test compound with 10 μ l of a substrate cocktail containing 500 μ g/ml peptide substrate and 0.2 mM ATP (+200 μ Ci/ml gamma-32P-ATP) in a 4x assay buffer. The reaction was initiated by the addition of 10 μ l of p38 kinase. Final assay conditions were 25 mM MOPS, pH
15 7.0, 26.25 mM beta-glycerol phosphate, 80 mM KCl, 22 mM $MgCl_2$, 3 mM $MgSO_4$, 1 mg/ml gelatin, 0.625 mM EGTA, 1 mM DTT, 125 μ g/ml peptide substrate, 50 μ M ATP, and 2.5 μ g/ml enzyme. After a 40 minute incubation at room temperature, the reaction was stopped by the addition of 10 μ l per reaction of 0.25 M phosphoric acid.

A portion of the reaction was spotted onto a disk of P81 phosphocellulose
20 paper, the filters were dried for 2 minutes and then washed 4x in 75 mM H_3PO_4 . The filters were rinsed briefly in 95% ethanol, dried, then placed in scintillation vials with liquid scintillation cocktail.

Alternatively, the substrate is previously biotinylated and the resulting reactions are spotted on SAM²TM streptavidin filter squares (Promega). The filters are
25 washed 4x in 2M NaCl, 4x in 2M NaCl with 1% phosphoric acid, 2x in water, and briefly in 95% ethanol. The filter squares are dried and placed in scintillation vials with liquid scintillation cocktail.

Counts incorporated are determined on a scintillation counter. Relative enzyme activity is calculated by subtracting background counts (counts measured in
30 the absence of enzyme) from each result, and comparing the resulting counts to those obtained in the absence of inhibitor. IC_{50} values were determined with curve-fitting

plots available with common software packages. Approximate IC_{50} values were calculated using formula

$$IC_{50} (app) = (A \times i)/(1-A)$$

where A = fractional activity and i = total inhibitor concentration.

5

Example 23

Activity of the Invention Compounds

The activity of the invention compounds was tested as described above. The compounds tested were 4-benzylpiperidinyl or 4-benzylpiperazinyl indole-5- or 6-carboxamides. In general, the piperidinyl derivative was superior to the
10 corresponding piperazinyl. The IC_{50} for inhibition of p38 α is shown in Table 1.

Table 1		
Ring position	Piperidinyl IC_{50} μM	Piperazinyl IC_{50} μM
5	0.150, 0.242	1.71, 1.78
6	0.462, 0.462	5.52, 4.97
3	0.2	5.44
4	0.2	1.55
2	3.26	>30
7	>30	>30

It is also seen that positioning the piperidinyl or piperazinyl substituent in positions 3, 4, 5 and 6 leads to greater activity than positioning the substituents in
15 positions 2 or 7.

The same compounds were tested for their specificity for p38 α as compared to p38 β . The results are shown in Table 2.

Table 2						
Ring position	Piperidinyl			Piperazinyl		
	P38- β IC_{50} μM	p38 α IC_{50} μM	IC_{50} ratio β/α	P38- β IC_{50} μM	p38- α IC_{50} μM	IC_{50} ratio β/α
5	3.02	0.150	20.1	25.8	1.71	15.1
6	3.83	0.462	8.27	39.1	5.52	7.08

Activity with regard to p38 β was also tested for compounds of the invention to determine the influence of the position of the piperidinyl or piperazinyl substituent. The influence of substitution on the benzyl moiety attached to the 4-position of the piperazine or piperidine was also tested. The results are shown in Table 3 in terms of the percent inhibition of p38 β activity at 50 μ M concentration of the compound.

Table 3			
Ring position	Benzyl substitution	Piperidinyl	Piperazinyl
5	-	96	59
6	-	92	56
3	-	96	77
4	-	96	68
2	-	12	27
7	-	45	7
5	4-Cl		77

Substituting benzimidazole for the indole moiety in the compounds of the invention also resulted in significant inhibition of p38 β when the compounds were tested at 50 μ M. 4-Benzyl piperidinyl benzimidazole-5-carboximide showed 85% inhibition; 4-(3-chlorobenzyl)piperizinyll benzimidazole-5-carboxamide showed 66% inhibition.

The compounds of the invention, generally, are specific for p38 α as compared to p38- β : It is seen that the specificity for α as opposed to β is generally of the order of ten-fold.

The specificity of the compounds of the invention was also tested with respect to other kinases, including p38- γ , ERK-2, PKA, PKC, cdc-2, EGF-R, and DNA-PK as shown in Table 4. The compounds tested are the 4-benzylpiperidinyl indole-5- and 6-carboxamides with the number indicating the ring position of the carboxamide.


Table 4		
KINASE	IC ₅₀ μ M 5	IC ₅₀ μ M 6
p38- α	0.150	0.462
p38- γ	228	>300
ERK-2	>300	>300
PKA	430	>500
PKC	>500	>500
cdc2	>500	>500
EGF-R	>500	>500
DNA-PK	>500	450

The results are given in terms of approximate IC₅₀ (μ M) values when the compounds were tested at 50 μ M and calculated using the formula in Example 22.


The exception is for p38 α values where the IC₅₀ values were determined from concentration dependent curve fitting analysis.

As shown, all of the compounds tested are highly specific for p38 α as compared to these additional kinases.





Table 5 shows the inhibition of p38 α by invention compounds that are 4-(benzyl-piperidinyl)indole-5-carboxamides or 4-[(4-fluorobenzyl) piperidinyl]indole-5-carboxamides, i.e., compounds of formulas (1) or (2):



wherein R¹ is of formula (11) or (12):




The values given are IC₅₀ in μ M.


Table 5						
Formula	R ³	R ¹	R ²	Z ¹	Z ²	IC ₅₀
1	H	(11)	H	N	CH	0.635
1	H	(11)	H	CH	CH	0.523, 0.577
2	CH ₃ O	(12)	H	CH	CH	0.159
2	Cl	(11)	Et ₂ N(CH ₂) ₃ -	CH	CH	0.199
1	H	(12)	Et ₂ N(CH ₂) ₂ -	CH	CH	0.354
2	CH ₃ O	(11)	H	CCONH(CH ₂) ₂ NMe ₂	CH	0.0646
1	H	(12)	H	N	CH	0.39
1	H	(11)	H	CCH ₂ - 	CH	6.57
1	H	(11)		CH	CH	0.871
1	H	(11)		CH	CH	0.405
1	H	(11)		CH	CH	0.8
1	Cl	(11)	H	CH	CCH ₃	0.489
1	H	(11)	Et ₂ N(CH ₂) ₃ -	CH	CH	0.474
2	MeO	(11)	Et ₂ N(CH ₂) ₂ -	CH	CH	0.184
1	Cl	(11)	H	CH	CCH ₃	(43% at 0.2 μM)

Several other compounds of the invention were also tested. A compound of formula (3) -- i.e., where the carboximide is in the 6-position, R¹ is of formula (11), R³ is H, R² is H, Z¹ is CCOCF₃ and Z² is CH showed 41% inhibition at 1 μM. Similarly,

5 a compound where R¹ is of formula (11) and substituted at the 6-position of the

indole, R² is  and both Z¹ and Z² are CH, showed an IC₅₀ of 0.505 μM.


Two compounds where R¹ is of the formula:



were also tested. In one, of formula (2) wherein R³ is MeO, R² is H and both Z¹ and Z² are CH, gave 63% inhibition at 0.2 μM; in the other case, which is of formula (3)

10


wherein R^3 is H, R^2 is H, Z^1 is N and Z^2 is CH, the IC_{50} was 2.15 μ M. Finally, one compound was tested wherein R^1 was of the formula:



and wherein the invention compound is of formula (3) wherein R^3 is H, R^2 is H, and
5 both Z^1 and Z^2 are CH, gave 51% inhibition at 1 μ M.

Claims

1. A compound of the formula:




and the pharmaceutically acceptable salts thereof,

- 5 wherein each of Z^1 and Z^2 is independently CR^4 or N;

- where each R^4 is independently H or is alkyl (1-6C) or aryl, each of said alkyl or aryl optionally including one or more heteroatoms selected from O, S and N and optionally substituted by one or more of halo, OR, SR, NR_2 , RCO, COOR, $CONR_2$, OOCR, or $NROCR$ where R is H or alkyl (1-6C), or by one or more CN or =O, or by one or more aliphatic or aromatic 5- or 6-membered rings optionally containing 1-2 heteroatoms;
- 10

R^1 is



wherein

- 15 X^1 is CO or an isostere thereof;

m is 0 or 1;

Y is optionally substituted alkyl, optionally substituted aryl, or optionally substituted arylalkyl or two Y taken together may form an alkylene (2-3C) bridge;

n is 0, 1 or 2;

- 20 Z^3 is CH or N;

X^2 is CH, CH_2 or an isostere thereof; and

Ar consists of one or two phenyl moieties directly coupled to X^2 optionally substituted by halo, nitro, alkyl (1-6C), alkenyl (1-6C), alkynyl (1-6C), CN or CF_3 , or


by RCO, COOR, CONR₂, NR₂, OR, SR, OOCR or NROCR wherein R is H or alkyl (1-6C) or by phenyl, itself optionally substituted by the foregoing substituents;

R² is H, or is alkyl (1-6C) or aryl, each of said alkyl or aryl optionally including one heteroatom which is O, S or N, and optionally substituted by one or more of halo, OR, SR, NR₂, RCO, COOR, CONR₂, OOCR, or NROCR where R is H or alkyl (1-6C), or by one or more CN or =O, or by one or more aliphatic or aromatic 5- or 6-membered rings optionally containing 1-2 heteroatoms;

R³ is H, halo, NO₂, alkyl (1-6C), alkenyl (1-6C), alkynyl (1-6C), CN, OR, SR, NR₂, RCO, COOR, CONR₂, OOCR, or NROCR where R is H or alkyl (1-6C).

10

2. The compound of claim 1 which is of the formula



3. The compound of claim 1 wherein m is 1 and wherein n is 0.

15



4. The compound of claim 1 wherein X¹ is CO.

5. The compound of claim 1 wherein X² is CH₂.



20 CH₂.

6. The compound of claim 2 wherein n is 0, m is 1, X¹ is CO and X² is

7. The compound of claim 1 wherein Z¹ and Z² are CR⁴.

8. The compound of claim 6 wherein Z^1 and Z^2 are CR^4 .
9. The compound of claim 1 wherein Z^1 is N and Z^2 is CH.
- 5 10. The compound of claim 6 wherein Z^1 is N and Z^2 is CH.
11. The compound of claim 2 which is of the formula (2).
12. The compound of claim 6 which is of the formula (2).
- 10 13. The compound of claim 2 wherein R^3 is halo or OR where R is alkyl (1-6C).
14. The compound of claim 6 wherein R^3 is halo or OR where R is alkyl
15 (1-6C).
15. The compound of claim 1 wherein Z^3 is CH.
16. The compound of claim 6 wherein Z^3 is CH.
- 20 17. The compound of claim 1 wherein Ar is  wherein each X^3 is independently alkyl (1-6C), halo, OR, or NR_2 and p is 0, 1, 2 or 3.
- 25 18. The compound of claim 6 wherein Ar is  wherein each X^3 is independently alkyl (1-6C), halo, OR, or NR_2 and p is 0, 1, 2 or 3.

19. The compound of claim 6 which is of the formula:



or having the structure of formula (5) or (6) wherein the positions on the indole or benzimidazole nucleus occupied by R³ and the substituent illustrated as R¹ are reversed,

5 wherein R², R³ and R⁴ are as defined in claim 1, and each X³ is independently halo, alkyl (1-6C), OR, or NR₂, wherein R is H or alkyl (1-6C) and p is 0, 1, 2 or 3.

20. The compound of claim 19 wherein p is 0 or p is 1 or 2 and each X³ is halo or OR where R is alkyl (1-3C).

10

21. The compound of claim 19 wherein R⁴ is H or is of the formula CONY' wherein Y' is alkyl, aryl or arylalkyl optionally containing one or two heteroatoms.

22. The compound of claim 19 wherein R² is H.

15

23. The compound of claim 19 wherein R³ is H, halo, or OR, wherein R is alkyl (1-6C).


24. The compound of claim 23 wherein R³ is chloro or methoxy.

20

25. The compound of claim 19 wherein the substituent R¹ shown in the 5-position of the indole or benzimidazole nucleus is at the 6-position and R³ is at the 5-position.

26. The compound of claim 19 wherein the substituents in formulas (5) and (6) are in these positions as shown.

27. The compound of claim 19 which is of the formula




wherein R^2 , R^4 , X^3 and p are as defined in claim 19.

28. The compound of claim 27 wherein at least one of R^2 and R^4 is a polar group.

29. The compound of claim 27 wherein R^4 is H or is of the formula $R_2N(CH_2)_nNHCO$ wherein n is an integer of 1-3 and each R is independently H or alkyl (1-6C) or wherein the R s taken together form a ring optionally containing a heteroatom which is S, O or N.

30. The compound of claim 19 which is of the formula



wherein R^2 , X^3 and p are as defined in claim 19.

31. The compound of claim 30 wherein R^2 is a polar group.

32. The compound of claim 29 wherein R^4 is H or is of the formula $R_2N(CH_2)_nNHCO$ wherein n is an integer of 1-3 and each R is independently H or alkyl (1-6C) or wherein the Rs taken together form a ring optionally containing a heteroatom which is S, O or N.

5

33. The compound of claim 1 which is

4-benzylpiperidinyl indole-5-carboxamide;

4-chloro-4-benzylpiperidinyl indole-5-carboxamide;

6-chloro-4-benzylpiperidinyl indole-5-carboxamide;

10 4-chloro-(4-(4-fluorobenzyl) piperidinyl)-indole-5-carboxamide;

6-chloro-(4-(4-fluorobenzyl) piperidinyl)-indole carboxamide;

4-methoxy-(4-benzylpiperidinyl)-indole-5-carboxamide;

6-methoxy-(4-benzylpiperidinyl)-indole-5-carboxamide;

4-methoxy-(4-(4-fluorobenzyl) piperidinyl)-indole-5-carboxamide;

15 6-methoxy-(4-(4-fluorobenzyl) piperidinyl)-indole-5-carboxamide;

N-(3-cyclohexylmethylamino-2-hydroxypropyl)-(4-benzylpiperidinyl)-indole-5-carboxamide;

N-(3-N-methylpiperazinyl-2-hydroxypropyl)-(4-benzylpiperidinyl)-indole-5-carboxamide;

20 N-(3-benzylamino-2-hydroxypropyl)-(4-benzylpiperidinyl)-indole-5-carboxamide;

N-[3-{(4-methoxybenzyl)-amino}-2-hydroxypropyl-](4-benzylpiperidinyl)-indole-5-carboxamide;

25 N-{3-n-propylamino-2-hydroxypropyl}-(4-benzylpiperidinyl)-indole-5-carboxamide;

N-(4-pyridoyl)-(4-benzylpiperidinyl)indole-5-carboxamide;

N-(4-pyridylmethyl)-(4-benzylpiperidinyl)-indole-5-carboxamide;

N-methylacetyl-(4-benzylpiperidinyl)-indole-5-carboxamide;

N-acetyl-4-benzylpiperidinyl indole-5-carboxamide;

30 N-(n-propylamide)acetyl 4-benzylpiperidinyl indole-5-carboxamide;

4-benzylpiperidinyl-indole-5-carboxamide-1-acetic acid-n-butylamide;

4-benzylpiperidiny-indole-5-carboxamide-1-acetic acid 4-methoxybenzyl
amide;

3-(2-methoxyethylaminocarboxamidyl)-(4-benzylpiperidiny)indole-5-
carboxamide;

5 3-(2-methylaminoethylaminocarboxamidyl)-(4-benzylpiperidiny)indole-5-
carboxamide;

3-(2-aminoethylaminocarboxamidyl)-(4-benzylpiperidiny)indole-5-
carboxamide;

10 3-(4-benzylpiperidinylcarboxamidyl)-(4-benzylpiperidiny)indole-5-
carboxamide;

3-(4-benzylpiperidinylcarboxamidyl)-(4-benzylpiperidiny)indole-6-
carboxamide;

3-(4-fluorobenzylcarboxamidyl)-(4-benzylpiperidiny)indole-5-carboxamide;

15 3-[2-(3,5-dimethoxyphenyl)ethylcarboxamidyl]-(4-benzylpiperidiny)indole-5-
carboxamide;

6-methoxy-(4-benzylpiperidiny)indole-5-carboxamide;

3-trifluoroacetyl-(4-benzylpiperidiny)indole-5-carboxamide;

6-methoxy-3-(2-dimethylaminoethylamino)carboxamidyl-(4-
benzylpiperidiny)indole-5-carboxamide;

20 3-trifluoroacetyl-4-benzylpiperidinylindole-5-carboxamide;

4-benzylpiperidiny indole-5-carboxamide-3-carboxylic acid;

3-(2-dimethylamino)ethylaminocarboxamidyl-(4-benzylpiperidiny)indole-5-
carboxamide;

or is a compound as set forth in Table 5.

25

34. The compound of claim 32 which is

4-benzylpiperdiny indole-5-carboxamide;


3-[2-dimethylaminoethylaminocarbonyl]-4-benzylpiperidiny-6-methoxy
indole-5-carboxamide; or

30 4-benzylpiperidiny-6-methoxy benzimidazole-5-carboxamide.

35. A method to treat a condition characterized by a proinflammation response which method comprises administering to a subject in need of such treatment a compound of claim 1 or a pharmaceutical composition thereof.

5 36. The method of claim 35 wherein said condition characterized by inflammation is acute respiratory distress syndrome, asthma, chronic obstructive pulmonary disease, uveitis, IBD, acute renal failure, head trauma, or ischemic/reperfusion injury.

10 37. A method to treat a heart condition associated with cardiac failure which method comprises administering to a subject in need of such treatment a compound of the formula



wherein R^1 , R^2 , R^3 , Z^1 , and Z^2 are as defined in claim 1, or administering a pharmaceutical composition thereof.

15

38. The method of claim 37 wherein said chronic heart condition is congestive heart failure, cardiomyopathy or myocarditis.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/11222

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D235/06 C07D401/06 C07D401/14 C07D209/08 C07D403/06
A61K31/445 A61K31/40 A61K31/41

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 431 945 A (MERCK & CO INC) 12 June 1991 (1991-06-12) claims; examples 24-26	1-4, 7, 11
X	EP 0 318 235 A (TAKEDA CHEMICAL INDUSTRIES LTD) 31 May 1989 (1989-05-31) see pages 21-22, working example 14 claims	1-4, 7, 11
X	EP 0 709 384 A (MERCK PATENT GMBH) 1 May 1996 (1996-05-01) claims	1-26
X	EP 0 831 090 A (ADIR) 25 March 1998 (1998-03-25) see page 3, lines 10-32 claims	1-26
	----- -/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

22 July 1999

Date of mailing of the international search report

29/07/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Henry, J

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/11222

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 06715 A (SMITHKLINE BEECHAM CORPORATION) 19 February 1998 (1998-02-19) cited in the application claims -----	1,35-38
P,X	WO 98 28292 A (SMITHKLINE BEECHAM CORPORATION) 2 July 1998 (1998-07-02) cited in the application claims -----	1,35-38

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/ 11222

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claim(s) 35-38
is(are) directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims: it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/11222

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0431945	A	12-06-1991	US 5032598 A	16-07-1991
			CA 2031693 A	09-06-1991
			JP 3181461 A	07-08-1991
			US 5215989 A	01-06-1993
EP 0318235	A	31-05-1989	JP 1230570 A	14-09-1989
			US 4937246 A	26-06-1990
EP 0709384	A	01-05-1996	DE 4438810 A	02-05-1996
			DE 19526269 A	23-01-1997
			AT 174915 T	15-01-1999
			AU 703487 B	25-03-1999
			AU 3443595 A	09-05-1996
			BR 9505008 A	14-10-1997
			CA 2161618 A	01-05-1996
			CN 1128762 A	14-08-1996
			CZ 9502816 A	15-05-1996
			DE 59504622 D	04-02-1999
			ES 2128629 T	16-05-1999
			FI 955184 A	01-05-1996
			HU 76068 A	30-06-1997
			JP 8225569 A	03-09-1996
			NO 954349 A	02-05-1996
			PL 311185 A	13-05-1996
			SK 135595 A	05-06-1996
			TR 960366 A	21-06-1996
			US 5698553 A	16-12-1997
			ZA 9509170 A	27-05-1996
EP 0831090	A	25-03-1998	FR 2753706 A	27-03-1998
			AU 3836297 A	26-03-1998
			BR 9704757 A	01-09-1998
			CA 2216664 A	20-03-1998
			HU 9701561 A	28-05-1998
			JP 10101645 A	21-04-1998
			NO 974253 A	23-03-1998
			NZ 328789 A	26-02-1998
			PL 322192 A	30-03-1998
WO 9806715	A	19-02-1998	EP 0922042 A	16-06-1999
WO 9828292	A	02-07-1998	NONE	